



This is an official
CDC Health Update

Distributed via Health Alert Network
March 5, 2013, 1:45PM
10281-CHU-03-05-2013-NECC

Notice to Clinicians: Continued Vigilance Urged for Fungal Infections among Patients Who Received Contaminated Steroid Injections

Summary


CDC continues to receive new reports of fungal infection among patients who were given injections of contaminated methylprednisolone acetate (MPA¹) from the New England Compounding Center (NECC) in Framingham, Mass. Most of these recent cases have been localized spinal or paraspinal infections (e.g., epidural abscesses) in patients, although new cases of meningitis or arachnoiditis also have been reported. Because many of these new cases are among patients with minimal symptoms, CDC is re-emphasizing the recommendation for clinicians to remain vigilant for fungal infections, especially in patients with mild or even baseline symptoms, and consider evaluation with magnetic resonance imaging (MRI) if clinically warranted. This Health Alert Network (HAN) notice provides the following:

- Information about the current status of the outbreak;
- Recommendations for clinical management and follow-up of exposed patients;
- Information about new revisions to web-based interim clinical guidance (http://www.cdc.gov/hai/outbreaks/clinicians/guidance_cns.html); and
- Notice of an upcoming CDC conference call to provide clinicians with additional diagnostic and treatment information.

Status of Fungal Disease Outbreak

As of March 4, 2013, a total of 720 cases, which includes 48 deaths, have been reported in 20 states. Current information about the outbreak, including case counts and distribution by state, and clinician and patient guidance, is available online at <http://www.cdc.gov/hai/outbreaks/meningitis.html>.

Fungal meningitis, often with a mild clinical presentation, was the predominant clinical syndrome reported among case-patients during the first several weeks of the outbreak (figure). Over the past several months, there has been a marked decrease in reports of fungal meningitis, but CDC continues to receive reports of localized spinal and paraspinal infections, which include epidural abscess, phlegmon, arachnoiditis, and discitis. Additionally, some of these newly identified case-patients had initially tested negative for signs of a fungal infection (either by lumbar puncture or MRI) and have subsequently developed fungal infection, indicating a prolonged incubation period.



After the recall of NECC steroid medications on September 26, state and local health departments identified almost 14,000 people in 23 states who were potentially exposed to the implicated MPA; of these, an estimated 11,000 individuals received spinal or paraspinal injections. Through active notification by clinics with assistance from states and CDC in early October, nearly all of these exposed persons were contacted at least once and informed of their risk for fungal infection as a result of receiving injections with contaminated medication.

Despite this and subsequent patient outreach efforts, CDC and public health partners remain concerned about the potential for some exposed patients to have localized fungal infections that have gone unrecognized. These infections may be unrecognized because some patients have not continued to receive close clinical follow-up or because they have not recognized symptoms suggestive of a localized infection, which may be difficult to distinguish from their baseline chronic pain.

As described in CDC's HAN update on December 20 (<http://emergency.cdc.gov/HAN/han00338.asp>), MRI testing was done on 128 patients in Michigan, Tennessee, and North Carolina who had no previous evidence of infection and had new or worsening symptoms at or near the site of their spinal or paraspinal injection. Of these, 67 (52%) had findings suggestive of localized infection. In addition, of 109 different patients reporting persistent but baseline symptoms at or near the site of their spinal or paraspinal injection, 15 (14%) also had abnormal MRI findings suggestive of infection, and 27 (25%) had non-specific enhancement of soft tissue or other paraspinal structures. The clinical significance of these findings is unclear; however, there is a theoretical risk that failure to diagnose these infections in a timely fashion could result in poor outcomes for patients (e.g., neurologic compromise, osteomyelitis, or progression to meningitis).

Patient and Clinician Recommendations

Early in the outbreak, CDC advised clinicians to closely monitor and evaluate patients who received injections of implicated MPA. Additional guidance was provided in HAN updates issued on November 20 (<http://emergency.cdc.gov/HAN/han00335.asp>) and December 20 (<http://emergency.cdc.gov/HAN/han00338.asp>). Because of the possibility that some patients may have unrecognized, localized fungal infections, CDC is re-emphasizing the following recommendations for patients who received a spinal or paraspinal injection with implicated MPA:

Patients

Patients who received an injection in or near their spine from one of the three implicated lots of MPA¹ and who have any symptoms at or near the site of their injection should seek evaluation by their medical provider for the possibility of a localized infection, such as an epidural abscess. This includes patients who initially received steroid injections for pain and continue to have persistent baseline pain.

Clinicians

As a part of continued monitoring of patients who received an injection with implicated MPA, clinicians should consider re-evaluating patients who received a spinal or paraspinal injection with implicated MPA for signs and symptoms suggestive of infection, including any symptoms at or near the site of their injection. Because of the prolonged incubation period for these infections, this guidance pertains both to patients who have not been previously evaluated and to those who have already had a prior negative evaluation (e.g., normal cerebrospinal fluid profile, normal findings on MRI) but continue to have complaints:

- In patients with new or worsening symptoms at or near the site of their injection, clinicians should obtain an MRI with contrast of the symptomatic area(s).
- In patients with persistent but baseline symptoms, clinicians should consider obtaining an MRI with contrast of the symptomatic area(s) because the presentation of spinal or paraspinal infections can be subtle, and may be difficult to distinguish from a patient's baseline chronic pain.
- In some cases, radiologic evidence of abscess or phlegmon has become apparent on repeat MRI studies performed subsequent to an initially normal imaging procedure. Clinicians should therefore have a low threshold for repeat MRI studies in patients who continue to have symptoms localizing to the site of injection, even after a normal study. However, the optimal duration between MRI studies is unknown.
- Clinicians should also consider reviewing MRI results with a neuroradiologist because of potential difficulties in interpreting imaging results for these patients.

Revised Clinical Guidance and Clinician Information Call

In response to input from expert consultants on fungal disease and physicians who have been treating patients affected by this outbreak, CDC has revised its Interim Treatment and Diagnostic Guidance for Central Nervous System and Parameningeal Infections Associated with Injection of Contaminated Steroid Products (http://www.cdc.gov/hai/outbreaks/clinicians/guidance_cns.html). The revisions include addition of new information on several topics, including:

- Surgical management of parameningeal disease
- Duration of antifungal treatment
- Monitoring clinical status after cessation of antifungal treatment
- Information on non-first-line medications (e.g., posaconazole or itraconazole)

A conference call for clinicians interested in obtaining additional information about the management and treatment of patients with fungal illness associated with this outbreak has been scheduled for March 13 at 5:00 p.m. The presenter will be Tom Chiller, M.D., medical officer, CDC. Registration and call-in information and other details about the conference call will be available on CDC's website <http://www.cdc.gov/hai/outbreaks/clinicians/index.html>.

NECC lots of methylprednisolone acetate (PF) 80mg/ml:

Methylprednisolone Acetate (PF) 80 mg/ml Injection, Lot #05212012@68, BUD 11/17/2012

Methylprednisolone Acetate (PF) 80 mg/ml Injection, Lot #06292012@26, BUD 12/26/2012

Methylprednisolone Acetate (PF) 80 mg/ml Injection, Lot #08102012@51, BUD 2/6/2013

DHEC contact information for reportable diseases and reporting requirements

Reporting of diseases is consistent with South Carolina Law requiring the reporting of diseases and conditions to your state or local public health department. (State Law # 44-29-10 and Regulation # 61-20) as per the DHEC 2013 List of Reportable Conditions available at:

<http://www.scdhec.gov/health/disease/reportables.htm>

Federal HIPAA legislation allows disclosure of protected health information, without consent of the individual, to public health authorities to collect and receive such information for the purpose of preventing or controlling disease. (HIPAA 45 CFR §164.512).

Regional Public Health Offices – 2013

Mail or call reports to the Epidemiology Office in each Public Health Region.

Region 1

Anderson, Oconee

220 McGee Road
Anderson, SC 29625
Phone: (864) 260-4358
Fax: (864) 260-5623
Nights / Weekends: 1-866-298-4442

Abbeville, Edgefield, Greenwood, Laurens, McCormick, Saluda

1736 S. Main Street
Greenwood, SC 29646
Phone: 1-888-218-5475
Fax: (864) 942-3690
Nights / Weekends: 1-800-420-1915

Region 2

Greenville, Pickens

PO Box 2507
200 University Ridge
Greenville, SC 29602-2507
Phone: (864) 372-3133
Fax: (864) 282-4373
Nights / Weekends: (864) 809-3825

Cherokee, Spartanburg, Union

PO Box 4217
151 E. Wood Street
Spartanburg, SC 29305-4217
Phone: (864) 372-3133
Fax: (864) 282-4373
Nights / Weekends: (864) 809-3825

Region 3

Chester, Lancaster, York

PO Box 817
1833 Pageland Highway
Lancaster, SC 29720
Phone: (803) 286-9948
Fax: (803) 286-5418
Nights / Weekends: 1-866-867-3886

Region 3 (continued)

Fairfield, Lexington, Newberry, Richland

2000 Hampton Street
Columbia, SC 29204
Phone: (803) 576-2749
Fax: (803) 576-2993
Nights / Weekends: 1-888-554-9915

Region 4

Chesterfield, Darlington, Dillon, Florence, Marlboro, Marion

145 E. Cheves Street
Florence, SC 29506
Phone: (843) 661-4830
Fax: (843) 661-4859
Nights / Weekends: (843) 601-7051

Clarendon, Kershaw, Lee, Sumter

PO Box 1628
105 North Magnolia Street
Sumter, SC 29150
Phone: (803) 773-5511
Fax: (803) 775-9941
Nights/Weekends: 803-458-1847

Region 5

Bamberg, Calhoun, Orangeburg

PO Box 1126
1550 Carolina Avenue
Orangeburg, SC 29116
Phone: (803) 268-5833
Fax: (803) 533-7134
Nights / Weekends: (803) 535-9265

Aiken, Allendale, Barnwell

222 Beaufort Street, NE
Aiken, SC 29801
Phone: (803) 642-1618
Fax: (803) 643-8386
Nights / Weekends: (803) 645-8167

Region 6

Georgetown, Horry, Williamsburg

1931 Industrial Park Road
Conway, SC 29526-5482
Phone: (843) 915-8804
Fax: (843) 365-0085
Nights/Weekends: (843) 340-4754

Region 7

Berkeley, Charleston, Dorchester

4050 Bridge View Drive, Suite 600
N. Charleston, SC 29405
Phone: (843) 953-0047
Fax: (843) 953-0051
Nights / Weekends: (843) 219-8470

Region 8

Beaufort, Colleton, Hampton, Jasper

219 S. Lemacks Street
Walterboro, SC 29488
Phone: (843) 441-1091
Fax: (843) 549-6845
Nights / Weekends: 1-843-441-1091

DHEC Bureau of Disease Control

Division of Acute Disease Epidemiology

1751 Calhoun Street
Box 101106
Columbia, SC 29211
Phone: (803) 898-0861
Fax: (803) 898-0897
Nights / Weekends: 1-888-847-0902



www.scdhec.gov

Categories of Health Alert messages:

Health Alert

Conveys the highest level of importance; warrants immediate action or attention.

Health Advisory

Provides important information for a specific incident or situation; may not require immediate action.

Health Update

Provides updated information regarding an incident or situation; unlikely to require immediate action.